

## REVIEW ARTICLE

# The Endocannabinoid System May Modulate Sleep Disorders in Aging

Eric Murillo-Rodríguez<sup>1,2,\*</sup>, Henning Budde<sup>2,3</sup>, André Barciela Veras<sup>2,4</sup>, Nuno Barbosa Rocha<sup>2,5</sup>, Diogo Telles-Correia<sup>2,6</sup>, Diogo Monteiro<sup>2,7,8</sup>, Luis Cid<sup>2,7,8</sup>, Tetsuya Yamamoto<sup>2,9</sup>, Sérgio Machado<sup>2,10</sup> and Pablo Torterolo<sup>2,11</sup>

<sup>1</sup>Laboratorio de Neurociencias Moleculares e Integrativas, Escuela de Medicina, División Ciencias de la Salud, Universidad Anáhuac Mayab, Mérida, Yucatán, México; <sup>2</sup>Intercontinental Neuroscience Research Group; <sup>3</sup>Faculty of Human Sciences, Medical School Hamburg, Hamburg, Germany; <sup>4</sup>Dom Bosco Catholic University, Campo Grande, Mato Grosso do Sul, Brazil; <sup>5</sup>School of Health, Polytechnic Institute of Porto, Porto, Portugal; <sup>6</sup>University of Lisbon, Faculty of Medicine, Lisbon, Portugal; <sup>7</sup>Sport Science School of Rio Maior-Polytechnic Institute of Santarém, Rio Maior, Portugal; <sup>8</sup>Research Center in Sport, Health and Human Development-CIDESD, Vila Real, Portugal; <sup>9</sup>Graduate School of Technology, Industrial and Social Sciences, Tokushima University, Tokushima, Japan; <sup>10</sup>Laboratory of Physical Activity Neuroscience, Physical Activity Sciences Postgraduate Program, Salgado de Oliveira University, Niterói, Brazil; <sup>11</sup>Laboratorio de Neurobiología del Sueño, Departamento de Fisiología, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

## ARTICLE HISTORY

Received: March 28, 2019

Revised: July 15, 2019

Accepted: July 17, 2019

DOI:

10.2174/1570159X17666190801155922

**Abstract:** Aging is an inevitable process that involves changes **across** life in multiple neurochemical, neuroanatomical, hormonal systems, and many others. In addition, these biological modifications lead to an increase in age-related sickness such as cardiovascular diseases, osteoporosis, neurodegenerative disorders, and sleep disturbances, among others that affect activities of daily life. Demographic projections have demonstrated that aging will increase its worldwide rate in the coming years. The research on chronic diseases of the elderly is important to gain insights into this growing global burden. Novel therapeutic approaches aimed for treatment of age-related pathologies have included the endocannabinoid system as an effective **tool** since this biological system shows beneficial effects in preclinical models. However, and despite these advances, little has been addressed in the arena of the endocannabinoid system as an option for treating sleep disorders in aging since experimental evidence suggests that some elements of the endocannabinoid system modulate the sleep-wake cycle. This article addresses this less-studied field, focusing on the likely perspective of the implication of the endocannabinoid system in the regulation of sleep problems reported in the aged. We conclude that beneficial effects regarding the putative efficacy of the endocannabinoid system as therapeutic tools in aging is either inconclusive or still missing.

**Keywords:** Cannabidiol, childhood, pharmacology, sleep.

## 1. INTRODUCTION

The demographic and epidemiologic projections have indicated growth in longevity worldwide in the coming years [1-3]. A critical distinction is between 1900, when only 4.1% of the 76 million people in the United States were  $\geq 65$  years old, 1950 with 8% of the total population under the same age, and 2000 with a percentage that had increased to 12.6% [4]. These numbers are enhancing worldwide since fertility and mortality rates are declining in most countries [5, 6]. Aging is also linked with the coexistence of two or more

chronic medical conditions, known as multimorbidity [7-9] leading to interactions between multiple disorders as well as among treatments prescribed for different disorders [9-12]. The geriatric-related health issues comprise multiple medical conditions such as hypertension, stroke, sleep disorders, among others [6, 13-18].

## 2. SLEEP MEDICINE AND AGING

The sleep-wake cycle is a complex biological phenomenon that involves the interaction of multiple complex neurobiological networks, including neuroanatomical, neurochemical, and genetic factors [19-24]. Normal sleep comprises two states, Non-REM (NREM) sleep, that has three phases (N1 to N3), and Rapid Eye Movement sleep (REMS). However, during aging, sleep displays aberrant features [25-29]. The sleep disorders have been classified by the Ameri-

\*Address correspondence to this author at the Laboratorio de Neurociencias Moleculares e Integrativas, Escuela de Medicina, División Ciencias de la Salud, Universidad Anáhuac Mayab, Carretera Mérida-Progreso Km. 15.5; A.P. 96 Cordemex C.P. 97310, Mérida, Yucatán, México; Tel: + 52 (999) 942-4800 Ext. 664; E-mail: [eric.murillo@anahuac.mx](mailto:eric.murillo@anahuac.mx)

can Academy of Sleep Medicine into the “International Classification of Sleep Disorders -third edition” (ICSD-3). Amongst the most common sleep complaints in the aged population are insomnia and excessive daytime sleepiness (EDS) or hypersomnolence [30-37].

Insomnia has been classified in several subtypes by polysomnographic studies and additional criteria. Sleep onset insomnia (persistent difficulty in sleep initiation), sleep maintenance insomnia (difficulty to maintain sleep during nighttime), early morning insomnia (identified with early morning awakenings), and psychophysiological insomnia (cognitive and behavioral elements, stress, intrusive thoughts, and attention bias) are reported in aging.

Based on the duration, insomnia is categorized in transient (few insomnia episodes before or during a stressful experience), short-term (events presented along a few weeks during an extended period of stress), or chronic (sleep disturbance and associated daytime symptoms have been present for at least three months) [30, 31, 36]. Remarkably, chronic insomnia affects 57% of the elderly in the USA [37-39].

On the other side of the coin, EDS is understood as the irresistible tendency to fall asleep during daytime [25, 27, 34, 40]. The diverse causes that originate EDS include, for instance, obstructive sleep apnea (interruptions in breathing during sleep for > 10 s leading to intermittent, partial (hypopnea), and/or complete (apnea) collapse of the upper airway during nighttime). Because of obstruction, any inspiratory effort causes snoring. In turn, this issue is commonly associated with the awakening of the subject across nighttime leading to sleep fragmentation. On the next day, and as a consequence, sleep fragmentation produces EDS [41-43], which is very **common** in older adults [44-47].

An additional sleep disturbance in aging is the periodic limb movement disorder (PLMD), characterized by repetitive cramping or jerking of the legs during sleep. PLMD is another cause of EDS [46-50]. Patients complaining **of** PLMD show involuntary and stereotypical movements of their limbs while they sleep. The movement of limbs across night induces recurrent awakenings that **limit** sleep consolidation and eventually, causes drowsiness on the next day. The prevalence of PLMD is higher among older adults [51-53].

Lastly, the neurodegeneration of a cluster of neurons located into the lateral hypothalamus, named orexin or hypocretin, seems to be the origin of the sleep disturbance named narcolepsy [54]. The core symptoms of this disease **comprise** extreme sleepiness (sleep attacks) during daytime and cataplexy [28, 54-56]. Although narcolepsy has been normally found during the adolescence, recent data have suggested the putative link between hypocretin/orexin and aging [57-61].

Currently, most of the sleep disturbances reported in aged subjects are managed by pharmacological interventions. Medical prescriptions for the elderly include antidepressants, antihistamines, sedative-hypnotics, among others [41, 49, 51, 53, 54, 62, 63]. However, since the study of the properties of the endocannabinoid system in modulating wider spectrum

of neurobiological phenomena, we would like to highlight the putative use of the endocannabinoid system elements in sleep disorders during aging.

### 3. THE ENDOCANNABINOID SYSTEM

During the 1990s, several experimental approaches were designed to describe the transmembrane proteins that pharmacologically responded to the principal compound of *Cannabis sativa*, the delta-9-tetrahydrocannabinol ( $\Delta^9$ -THC). At this date, it is widely accepted that the receptors that respond to  $\Delta^9$ -THC are CB<sub>1</sub> and CB<sub>2</sub> cannabinoid receptors [64-67]. In addition, the presence of both receptors has been mapped in different organs, including the human brain [68-71].

Next, the quest of the endogenous ligands that naturally bind to these receptors leads to the discovery of the first agonist to the cannabinoid receptors named arachidonylethanolamine or anandamide, and in later years, the second lipid with cannabimimetic properties was described: 2-arachidonylglycerol (2-AG) [72]. Currently, the family of the endocannabinoids comprises different compounds aside anandamide and 2-AG, such as virodhamine, noladin ether, and N-arachidonyldopamine [73-77]. In addition, the endocannabinoid system is integrated by several molecular elements, including the enzymes that synthesize/degrade anandamide (fatty acid hydrolase [FAAH]) or 2-AG (monoacylglycerol lipase [MAGL]), the anandamide membrane transporter (AMT) and the receptor channel TRPV1 (Transient Receptor Potential Vanilloid 1; [78-82]). Along the decades, preclinical studies have shown that the endocannabinoid system exerts critical neurobiological functions, some of them, with clinical relevance, including the control of the sleep-wake cycle [80, 83-93].

### 4. THE ENDOCANNABINOID SYSTEM IN HEALTH ISSUES

Multiple experimental models strongly suggest that the endocannabinoid system plays a critical role in numerous pathologies. Converging preclinical experiments, including a wider spectrum of data, from modulating the endogenous tone of anandamide, to the treatment with a direct CB<sub>1</sub> cannabinoid receptor agonist/antagonist, the pharmacological inhibition of the FAAH, or the genetic deletion of cannabinoid receptors, have been shown to control adverse medical conditions, such as schizophrenia, and ataxias. Amongst these pathological issues, important advances have been achieved in neurodegenerative disorders, including Parkinson's and Alzheimer disease [94-101]. Yet the hypothesis that the endocannabinoid system might be critically involved in ongoing pathologies associated with aging, such as sleep disorders, has been dismissed. However, an accumulating body of evidence is providing pieces to the puzzle to appreciate the putative role of the endocannabinoid system in the modulation of sleep disorders in the elderly.

### 5. AGING AND THE ENDOCANNABINOID SYSTEM

It is often argued that aging is associated with multiple deteriorating medical conditions, such as neurodegenerative disorders, leading to the dysregulation of endogenous neurochemical compounds. In this regard, the levels of anan-

damide and 2-AG have been described increased in Parkinson's disease patients compared to respective healthy subjects [102, 103]. Complementary, it is worth to emphasize that **in** patients with Alzheimer's disease, the FAAH gene expression has been found higher compared to age-matched controls [104-106]. Furthermore, the **lack** of the CB<sub>2</sub> cannabinoid receptors has been linked with schizophrenia-like pathologies whereas contents of anandamide are significantly enhanced in schizophrenic patients compared to respective controls [107-110]. **Taken** together, these data oblige us to address an important question that remains to be answered: Are the health disturbances the cause or consequence of the disruption of the endocannabinoid system functioning? While pertinent experimental approaches are aimed to address this concern, the body of evidence suggests that there is a neurobiological role of the endocannabinoid system in aging processes, including perhaps, sleep disturbances.

## 6. THE ENDOCANNABINOID SYSTEM MODULATING THE SLEEP-WAKE CYCLE

Significant achievements have been obtained by describing the modulatory role of the endocannabinoid system in the sleep-wake cycle. Pharmacological evidence from experiments on central or peripheral administrations **has** shown that elements of the endocannabinoid family participate in sleep control. For example, pioneer studies by the laboratory of Gerard Le Fur [111], Vincent Santucci, and coworkers demonstrated that the systemic injections of the CB<sub>1</sub> cannabinoid receptor antagonist SR 141716A (0.1, 0.3, 1, 3 and 10 mg/Kg) increased the time spent in wakefulness (W) while decreased slow-wave sleep (SWS) and REMS whereas opposite results were described by our group in 1998 after icv injection of anandamide [112]. Remarkably, the sleep-inducing effects of anandamide were blocked by SR141716A [113].

Another element of the endocannabinoid system involved in sleep modulation includes the FAAH. This assumption is supported by studies in which administrations either icv or into the lateral hypothalamus of URB597, a relatively selective inhibitor of FAAH, enhanced W and decreased SWS during the lights-on period [114, 115]. Previous findings have indicated that injections of URB597 enhance waking and decrease sleep. An unexpected result since one might think that inhibition of FAAH should increase anandamide contents and then, facilitates sleep onset. However, data suggest that the effects on sleep caused by URB597 might be due to the accumulation of oleoylethanolamide (OEA) and/or palmitoylethanolamide (PEA) since FAAH also synthesizes both lipids in greater amounts compared to anandamide. In line with this idea, our group demonstrated that the administration of OEA or PEA **increases** wakefulness and reduces SWS and REM sleep mimicking URB597's effects [114, 115]. Lastly, OEA behaves as an agonist with high-affinity to the PPAR- $\alpha$  [116], raising the possibility that PPAR- $\alpha$  could be the mechanism of action of OEA to induce waking [117, 118]. Further studies have also shown that additional system are engaged in the pharmacological effects on sleep of URB597, OEA or PEA, including the dopaminergic neurotransmission [114, 115, 117, 118].

In another set of experiments in which rats received AMT inhibitors, VDM-11 or OMDM-2, REMS was increased [119, 120]. This data suggested that REMS promotion could be as a consequence of the enhancement in the endogenous contents of anandamide.

To elucidate the role of the CB<sub>1</sub> cannabinoid receptor, FAAH and AMT in sleep homeostasis, experimental animals were subjected to total sleep deprivation and they received before the sleep rebound period, systemic injection of either SR141716A, URB597, or VDM-11 (5, 10, 20mg/Kg; ip; separately each one). As expected, SR141716A and URB597 blocked the sleep rebound whereas VDM-11 exerted opposite effects [121].

Finally, what might be the role of the endocannabinoid system in sleep medicine? The fact that anandamide has been associated with sleep apnea [86, 122-124] points to the possibility that the CB<sub>1</sub> cannabinoid receptor, FAAH, MAGL, or AMT may be also related to sleep disorders and they could act as criteria for diagnostic sleep disturbances.

Thus, the information available allows us to identify the elements of the endocannabinoid family as candidates for sleep control. Nevertheless, there is a lack of data demonstrating whether the endocannabinoid system may control sleep disorders in aging since most of the current data has been mostly reported in young or adult animals.

## 7. DOES THE ENDOCANNABINOID SYSTEM MODULATE SLEEP DISORDERS IN THE ELDERLY?


There is a significant amount of experimental and pre-clinical evidence demonstrating that the endocannabinoid system is linked with neurobiological processes related to aging. For example, murine models lacking the CB<sub>1</sub> cannabinoid receptors display an early onset of learning disturbances linked with aging. These findings have contributed to the design of drugs aimed to manage age-related human pathologies *via* the endocannabinoid system elements, such as the CB<sub>1</sub> cannabinoid receptors, anandamide, FAAH, AMT, amongst others [124-132]. We would like to describe a hypothetical framework regarding the likely modulatory role of the endocannabinoid system on sleep disorders in the elderly (Fig. 1).

### 7.1. The CB<sub>1</sub> Cannabinoid Receptor

Anandamide and 2-AG **promote** sleep *via* activation of the CB<sub>1</sub> cannabinoid receptor [92, 112, 113]. The available literature has indicated that the CB<sub>1</sub> cannabinoid receptor in aged rats seems to be decreased [125, 133].

### 7.2. The Endogenous Tone of Anandamide or 2-AG

Diurnal variation of endocannabinoids has been detected in cerebrospinal fluid, pons, hippocampus, or hypothalamus in rats [134, 135]. For example, anandamide in CSF showed an enhancement during the lights-on period whereas decreased across the lights-off period in young rats whereas anandamide and 2-AG levels are decreased in aging [126, 136, 137].



**Fig. (1).** The endocannabinoid system in aging. The CB<sub>1</sub> cannabinoid receptor, anandamide, 2-AG, monoacylglycerol lipase (MAGL), anandamide membrane transporter (AMT), and fatty acid amide hydrolase (FAAH) belong to the endocannabinoid system. Briefly, anandamide is synthesized by the activity of FAAH in the postsynaptic terminal. Next, the AMT displaces anandamide into the extracellular space to binding to the CB<sub>1</sub> cannabinoid receptor. In addition, the 2-AG also binds to the CB<sub>1</sub> cannabinoid receptor at the presynaptic terminal. As mentioned in the text, anandamide, 2-AG, the CB<sub>1</sub> cannabinoid receptor, FAAH, and AMT participate in sleep modulation. As shown in the drawing, the CB<sub>1</sub> cannabinoid receptor, anandamide, and 2-AG levels have been found decreased in aging. However, the activity of FAAH, AMT, and MAGL across aging remains to be described.

### 7.3. FAAH and MAGL

It is widely accepted that anandamide is synthesized on demand and regulated by the activity of the FAAH whereas 2-AG is predominantly hydrolyzed to arachidonic acid and glycerol by MAGL [138]. Thus, the activity of these enzymes in the regulation of anandamide and 2-AG contents is critical. In this regard, Pascual *et al.* (2014) demonstrated that FAAH activity was decreased in the frontal cortex from patients with Alzheimer's disease [139] whereas the similar findings were reported for MAGL in Alzheimer's disease experimental murine models [140-142]. In line with this data, Piyanova *et al.* (2015) found an elevated MAGL activity during aging [143].

### 7.4. AMT

As mentioned previously, the AMT behaves as a carrier-mediated transport for anandamide across the neuronal membrane. Unfortunately, limited data are available regarding the role of AMT in aging. To our understanding, the only research that has covered this issue comes from Maccarrone *et al.* (2001) who reported that mice lacking the CB<sub>1</sub> cannabinoid receptor showed a significant increase in AMT with age, whereas minor changes were found in wild-type animals [142].

Since aging is characterized by the progressive impairment of diverse physiological functions leading to developing medical disturbances, including neurodegenerative dis-



eases, then the study of the likely interaction of the endocannabinoid system with the mechanisms linked to age-related disturbances represents an interesting route for testing novel therapeutics. Here, we propose a conceptual framework linking the endocannabinoid system and the control of sleep disturbances in aging. However, our hypothetical proposal requires further evaluation by, for example, describing the distribution in the central nervous system of the CB<sub>1</sub> cannabinoid receptor in aging and how they are altered under specific sleep disorders. Second, further evidence is required to address the modulation of the endocannabinoid system using experimental models of sleep disturbances targeting cannabinoid receptors with agonists or antagonists, increasing anandamide and 2-AG endogenous tone by the inhibition of FAAH or MAGL activity as well as the functioning of AMT. Lastly, translational studies should provide perspectives on how to bridge from basic science to human health for the developing endocannabinoid-based diagnostics and therapies for sleep disturbances in aging.

#### 8. CANNABINOIDS SEEM TO BE PROMISING...BUT WAIT! WHAT ABOUT THE LONG-TERM EFFECTS IN THE ELDERLY DUE TO THE STIMULATION BY CANNABINOIDS DURING CHILDHOOD?


$\Delta^9$ -THC and cannabidiol (CBD) are the major chemicals in *Cannabis sativa*. While the first compound has been related to neurobiological disturbances, the second one does not exert psychotropic effects. Thanks to the research during the decades of the 1980s and 1990s, it was possible the characterization of the membrane binding sites for  $\Delta^9$ -THC. Currently, it is accepted that exogenous and endogenous cannabinoids bind to the CB<sub>1</sub> and CB<sub>2</sub> cannabinoid receptors [68, 69, 71, 72]. Due to the description of the cannabinoid

receptors in human organs, including the central nervous system, the localization of these receptors correlated the behavioral effects and the administration of  $\Delta^9$ -THC [144-147].

Due to CBD not inducing undesirable effects, it has been considered as a promising compound in the last decades. In line with this, CBD seems to manage several medical conditions such as psychosis, inflammation, asthma, autism, and pediatric epilepsy [148]. In recent years, it has been suggested the clinical application of CBD for managing anxiety and sleep disturbances such as posttraumatic stress disorder [149-153]. Thus, as one can suspect, the accelerating acceptance of CBD-based products aimed for treating several illnesses should raise questions whether long-uses of CBD could induce adverse effects in aging.

The importance of discussing the putative long-term effects of CBD on sleep in aging lies in the evidence that shows the interaction among this compound and some elements of the endocannabinoid system. For example, current approaches have suggested that CBD behaves as a CB<sub>1</sub> cannabinoid receptor negative allosteric modulator [154-157]. Moreover, CBD inhibits FAAH activity and modulates anandamide and 2-AG levels [158]. Thus, it seems reasonable to suspect that long-term exposure to CBD during early years might induce sleep disturbances in aging.

To increase awareness in regards to putative effects derived from long-term uses of CBD, various reports have demonstrated that this phytocannabinoid engages multiple neurobiological systems [148], including the increase of the extracellular contents of acetylcholine, adenosine, dopamine, and serotonin [159-161]. Since it has not been described which of these neurobiological pathways could be responsible for the presumably therapeutic properties of CBD, then



**Fig. (2).** The likely long-term neurobiological effects in aging after chronic cannabidiol (CBD) usage during childhood. Due to the susceptibility of the brain in childhood, it is possible that long term prescription of CBD might cause effects in learning, memory or causing sleep disorders in aging.

multiple neurobiological routes might be engaged in the long-term prescription of this compound. Due to CBD tested exerting positive outcomes in epilepsy [162-165], then this phytocannabinoid has been also evaluated as a therapeutical option for managing schizophrenia, Alzheimer disease, autism, among others pathological conditions [148-150]. Therefore, the chronic use of CBD might cause undesirable effects in aging. To **sum** concerns, on-line availability and accessibility to CBD to products which **do** not fulfill FDA standards [166], could add healthy risks due to the chronic use of CBD. If used at juvenile ages, the neuronal development might be under the influence of CBD since the brain networks are still actively developing during adolescence [167]. Hence, the long-term use of CBD might induce neuromolecular dysfunctions if treatment extends from childhood to adulthood as previously suggested (Fig. 2; [168-171]).

## 9. THE LIMITATIONS OF CURRENT EVIDENCE

At this date, the evidence of long-term disturbances in aging by lasting uses of CBD across childhood is limited and restricts to draw solid conclusions. We have identified that limitations of available literature include the variability of the experimental designs used animal model/clinical condition (epilepsy, anxiety, depression, *etc*), dosage (*e.g.*, low vs. high), route of administration (central vs. peripheral), frequency of usage (chronic vs. acute), among other conditions [145, 148, 172]. Moreover, achieving longitudinal studies that show time-varying exposure related to neurobiological effects in chronically CBD-treated subjects will provide experimental evidence to translate to medical areas where uses of CBD seems to be common such as pediatric epilepsy.

## CONCLUSIONS

The increase in the number of aged people around the world and age-related health disturbances is a worldwide challenge. To note, the costs of dementia in the USA was more than US\$800 billion in 2015, corresponding to a 35% increase compared to 2010 [1-6]. Thus, aging phenomena should promote the development of new therapeutical approaches.

During aging, the sleep-wake cycle includes abnormalities classified as sleep disorders, such as insomnia, obstructive sleep apnea, and excessive somnolence, among many others. **Since** projective models have indicated that the number of the oldest population would be higher, then this situation will also represent a public health challenge to assist age-related health issues, including sleep disorders. Here, we integrated the current knowledge of some sleep disorders in aging. The endocannabinoid system and its relationship with aging were revised as well. Lastly, the hypothetical link between the endocannabinoid system and sleep disorders in aging were proposed.

Moving our analysis from the endocannabinoid system and the neuronal circuits engaged on the sleep-wake cycle regulation is beyond the aim of the current Review since further complexity has been recognized by the localization of the elements of the endocannabinoid system in multiple sleep-related brain areas as well as the neuronal networks that **underlie** the influence of the endocannabinoid system in

sleep control. Since the CB<sub>1</sub> cannabinoid receptor, anandamide, 2-AG, and FAAH have been identified in sleep brain areas such as the cerebral cortex, hypothalamus and pons [173, 174], then we can assume that the endocannabinoid system might modulate the sleep-wake cycle. However, clinical implications between the endocannabinoid system functioning and sleep disturbances in aging require further investigation. Moreover, future research should focus on identifying the mechanisms of action of the multiple elements of the endocannabinoid system in aging and sleep disturbances as well as the risks of neuromolecular durable effects in aging if the chronic prescription of CBD is provided across childhood.

Given the medical regulations in several countries allowing the medicinal uses of *Cannabis sativa* or compounds derived of cannabinoids, and the on-line accessibility to CBD, a diversity of uncontrolled and available CBD-containing products (*e.g.* oils, foods, drinks, gums, *etc.*) **on** the internet, which some of them do not meet the rigorous FDA standards [168] would increase the influx of subjects with acute illnesses searching purported medicinal products that could cause additional concerns than resolution of these. Thus, legitimate concerns **are raised** about the risks of neurobiological durable effects in adulthood if CBD is prescribed during juvenile ages. Moreover, there is no solid evidence in regards **to** each *Cannabis sativa* ingredient, or its mixture could exert positive outcomes for treating sleep disorders.

Although a plethora of pharmacological evidence of the endocannabinoid system have recently been suggested relevant for its likely sleep-modulatory properties in aging, the possibility that some beneficial outcomes for age-related sleep disturbances regulation could be achieved by indirect mechanism such as the control of pain, anxiety, depression, among other medical conditions [175-177] remains to be unclear at present.

## CONSENT FOR PUBLICATION

Not applicable.

## FUNDING

None.

## CONFLICT OF INTEREST

We declare no competing interests. This work was supported by The University of California Institute for Mexico and the United States (UC MEXUS) and Consejo Nacional de Ciencia y Tecnología (CONACyT; Grant# CN-17-19) and Escuela de Medicina, Universidad Anáhuac Mayab (Grant: PresInvEMR2017) given to E M-R.

## ACKNOWLEDGEMENTS

All authors contributed substantially to the article.

## REFERENCES

- [1] World Health Organization. Global Health and Aging. National Institute on Aging National Institutes of Health U.S. Department of Health and Human Services. USA. NIH. Publication no. 11-7737. 2011.

- [2] Beard, J.R.; Officer, A.; de Carvalho, I.A.; Sadana, R.; Pot, A.M.; Michel, J.P.; Lloyd-Sherlock, P.; Epping-Jordan, J.E.; Peeters, G.M.E.E.G.; Mahanani, W.R.; Thiyagarajan, J.A.; Chatterji, S. The World report on ageing and health: a policy framework for healthy ageing. *Lancet*, **2016**, 387(10033), 2145-2154. [http://dx.doi.org/10.1016/S0140-6736(15)00516-4] [PMID: 26520231]
- [3] Götmarm, F.; Cafaro, P.; O'Sullivan, J. Aging Human Populations: Good for Us, Good for the Earth. *Trends Ecol. Evol. (Amst.)*, **2018**, 33(11), 851-862. [http://dx.doi.org/10.1016/j.tree.2018.08.015] [PMID: 30340868]
- [4] Ferrucci, L.; Giallauria, F.; Guralnik, J.M. Epidemiology of aging. *Radiol. Clin. North Am.*, **2008**, 46(4), 643-652, v. [http://dx.doi.org/10.1016/j.rcl.2008.07.005] [PMID: 18922285]
- [5] Rohde, J.E. Far more than 7 billion. *Lancet Infect. Dis.*, **2012**, 12(1), 12. [http://dx.doi.org/10.1016/S1473-3099(11)70338-3] [PMID: 22192129]
- [6] Beard, J.R.; Officer, A.; de Carvalho, I.A.; Sadana, R.; Pot, A.M.; Michel, J.P.; Lloyd-Sherlock, P.; Epping-Jordan, J.E.; Peeters, G.M.E.E.G.; Mahanani, W.R.; Thiyagarajan, J.A.; Chatterji, S. The World report on ageing and health: a policy framework for healthy ageing. *Lancet*, **2016**, 387(10033), 2145-2154. [http://dx.doi.org/10.1016/S0140-6736(15)00516-4] [PMID: 26520231]
- [7] Tugwell, P.; Knottners, J.A. Multimorbidity and Comorbidity are now separate MESH headings. *J. Clin. Epidemiol.*, **2019**, 105, vi-viii. [http://dx.doi.org/10.1016/j.jclinepi.2018.11.019] [PMID: 30522770]
- [8] Mayor, S. Commonly used approach for multimorbidity fails to improve quality of life. *BMJ*, **2018**, 361, k287. [http://dx.doi.org/10.1136/bmj.k2875]
- [9] Vetrano, D.L.; Palmer, R.; Marengoni, A.; Marzetti, E.; Lattanzio, F.; Roller-Wirnsberger, K.; Lopez Samaniego, L.; Rodríguez-Mañas, L.; Bernabei, R.; Onder, G. Joint Action ADVANTAGE WP4 group. Frailty and multimorbidity: a systematic review and meta-analysis. *J. Gerontol. A Biol. Sci. Med. Sci.*, **2018**.
- [10] Tisminetzky, M.; Gurwitz, J.H.; Fan, D.; Reynolds, K.; Smith, D.H.; Magid, D.J.; Sung, S.H.; Murphy, T.E.; Goldberg, R.J.; Go, A.S. Multimorbidity Burden and Adverse Outcomes in a Community-Based Cohort of Adults with Heart Failure. *J. Am. Geriatr. Soc.*, **2018**, 66(12), 2305-2313. [http://dx.doi.org/10.1111/jgs.15590] [PMID: 30246862]
- [11] Calderón-Larrañaga, A.; Fratiglioni, L. Multimorbidity research at the crossroads: developing the scientific evidence for clinical practice and health policy. *J. Intern. Med.*, **2019**, 285(3), 251-254. [http://dx.doi.org/10.1111/joim.12872] [PMID: 30625511]
- [12] Nicholson, K.; Makovski, T.T.; Griffith, L.E.; Raina, P.; Stranges, S.; van den Akker, M. Multimorbidity and comorbidity revisited: refining the concepts for international health research. *J. Clin. Epidemiol.*, **2019**, 105, 142-146. [http://dx.doi.org/10.1016/j.jclinepi.2018.09.008] [PMID: 30253215]
- [13] Childs, B.G.; Durik, M.; Baker, D.J.; van Deursen, J.M. Cellular senescence in aging and age-related disease: from mechanisms to therapy. *Nat. Med.*, **2015**, 21(12), 1424-1435. [http://dx.doi.org/10.1038/nm.4000] [PMID: 26646499]
- [14] Fung, C.H.; Vitiello, M.V.; Alessi, C.A.; Kuchel, G.A. AGS/NIA Sleep Conference Planning Committee and Faculty. Report and Research Agenda of the American Geriatrics Society and National Institute on Aging Bedside-to-Bench Conference on Sleep, Circadian Rhythms, and Aging: New Avenues for Improving Brain Health, Physical Health, and Functioning. *J. Am. Geriatr. Soc.*, **2016**, 64(12), e238-e247. [http://dx.doi.org/10.1111/jgs.14493] [PMID: 27858974]
- [15] Miner, B.; Kryger, M.H. Sleep in the Aging Population. *Sleep Med. Clin.*, **2017**, 12(1), 31-38. [http://dx.doi.org/10.1016/j.jsmc.2016.10.008] [PMID: 28159095]
- [16] Mander, B.A.; Winer, J.R.; Walker, M.P. Sleep and Human Aging. *Neuron*, **2017**, 94(1), 19-36. [http://dx.doi.org/10.1016/j.neuron.2017.02.004] [PMID: 28384471]
- [17] Musiek, E.S.; Bhimasani, M.; Zangrilli, M.A.; Morris, J.C.; Holtzman, D.M.; Ju, Y.S. Circadian Rest-Activity Pattern Changes in Aging and Preclinical Alzheimer Disease. *JAMA Neurol.*, **2018**, 75(5), 582-590. [http://dx.doi.org/10.1001/jamaneurol.2017.4719] [PMID: 29379963]
- [18] Dong, X.; Milholland, B.; Vijg, J. Evidence for a limit to human lifespan. *Nature*, **2016**, 538(7624), 257-259. [http://dx.doi.org/10.1038/nature19793] [PMID: 27706136]
- [19] Murillo-Rodríguez, E.; Arias-Carrion, O.; Zavala-Garcia, A.; Sarro-Ramirez, A.; Huitron-Resendiz, S.; Arankowsky-Sandoval, G. Basic sleep mechanisms: an integrative review. *Cent. Nerv. Syst. Agents Med. Chem.*, **2012**, 12(1), 38-54. [http://dx.doi.org/10.2174/187152412800229107] [PMID: 22524274]
- [20] Spiegelhalter, K.; Riemann, D. Losing sleep. *Lancet Neurol.*, **2015**, 14(6), 571. [http://dx.doi.org/10.1016/S1474-4422(15)00065-4] [PMID: 25987277]
- [21] Arnulf, I. Sleep: what the day owes the night. *Lancet Neurol.*, **2015**, 14(1), 19-20. [http://dx.doi.org/10.1016/S1474-4422(14)70234-0] [PMID: 25496888]
- [22] Siclari, F.; Baird, B.; Perogamvros, L.; Bernardi, G.; LaRocque, J.J.; Riedner, B.; Boly, M.; Postle, B.R.; Tononi, G. The neural correlates of dreaming. *Nat. Neurosci.*, **2017**, 20(6), 872-878. [http://dx.doi.org/10.1038/nn.4545] [PMID: 28394322]
- [23] Hill, V.M.; O'Connor, R.M.; Shirasu-Hiza, M. Tired and stressed: Examining the need for sleep. *Eur. J. Neurosci.*, **2018**. [http://dx.doi.org/10.1111/ejn.14197] [PMID: 30295966]
- [24] Gent, T.C.; Bandarabadi, M.; Herrera, C.G.; Adamantidis, A.R. Thalamic dual control of sleep and wakefulness. *Nat. Neurosci.*, **2018**, 21(7), 974-984. [http://dx.doi.org/10.1038/s41593-018-0164-7] [PMID: 29892048]
- [25] Sateia, M.J. International classification of sleep disorders-third edition: highlights and modifications. *Chest*, **2014**, 146(5), 1387-1394. [http://dx.doi.org/10.1378/chest.14-0970] [PMID: 25367475]
- [26] Morin, C.M.; Drake, C.L.; Harvey, A.G.; Krystal, A.D.; Manber, R.; Riemann, D.; Spiegelhalter, K. Insomnia disorder. *Nat. Rev. Dis. Primers*, **2015**, 1, 15026. [http://dx.doi.org/10.1038/nrdp.2015.26] [PMID: 27189779]
- [27] Ruoff, C.; Rye, D. The ICSD-3 and DSM-5 guidelines for diagnosing narcolepsy: clinical relevance and practicality. *Curr. Med. Res. Opin.*, **2016**, 32(10), 1611-1622. [http://dx.doi.org/10.1080/03007995.2016.1208643] [PMID: 27359185]
- [28] Mahoney, C.E.; Cogswell, A.; Koralnik, I.J.; Scammell, T.E. The neurobiological basis of narcolepsy. *Nat. Rev. Neurosci.*, **2019**, 20(2), 83-93. [http://dx.doi.org/10.1038/s41583-018-0097-x] [PMID: 30546103]
- [29] Dauvilliers, Y.; Schenck, C.H.; Postuma, R.B.; Iranzo, A.; Luppi, P.H.; Plazzi, G.; Montplaisir, J.; Boeve, B. REM sleep behaviour disorder. *Nat. Rev. Dis. Primers*, **2018**, 4(1), 19. [http://dx.doi.org/10.1038/s41572-018-0016-5] [PMID: 30166532]
- [30] Jausent, I.; Bouyer, J.; Ancelin, M.L.; Akbaraly, T.; Pérès, K.; Ritchie, K.; Besset, A.; Dauvilliers, Y. Insomnia and daytime sleepiness are risk factors for depressive symptoms in the elderly. *Sleep (Basel)*, **2011**, 34(8), 1103-1110. [http://dx.doi.org/10.5665/SLEEP.1170] [PMID: 21804672]
- [31] Wennberg, A.M.; Canham, S.L.; Smith, M.T.; Spira, A.P. Optimizing sleep in older adults: treating insomnia. *Maturitas*, **2013**, 76(3), 247-252. [http://dx.doi.org/10.1016/j.maturitas.2013.05.007] [PMID: 23746664]
- [32] Akbaraly, T.N.; Jausent, I.; Besset, A.; Bertrand, M.; Barberger-Gateau, P.; Ritchie, K.; Ferrie, J.E.; Kivimaki, M.; Dauvilliers, Y. Sleep complaints and metabolic syndrome in an elderly population: the Three-City Study. *Am. J. Geriatr. Psychiatry*, **2015**, 23(8), 818-828. [http://dx.doi.org/10.1016/j.jagp.2014.10.001] [PMID: 25499672]
- [33] Mattis, J.; Sehgal, A. Circadian Rhythms, Sleep, and Disorders of Aging. *Trends Endocrinol. Metab.*, **2016**, 27(4), 192-203. [http://dx.doi.org/10.1016/j.tem.2016.02.003] [PMID: 26947521]
- [34] Gabelle, A.; Gutierrez, L.A.; Jausent, I.; Navucet, S.; Grasselli, C.; Bennis, K.; Marelli, C.; David, R.; Andrieu, S.; Berr, C.; Vellas, B.; Dauvilliers, Y. Excessive Sleepiness and Longer Nighttime in

- Bed Increase the Risk of Cognitive Decline in Frail Elderly Subjects: The MAPT-Sleep Study. *Front. Aging Neurosci.*, **2017**, 9, 312.  
[http://dx.doi.org/10.3389/fnagi.2017.00312] [PMID: 29033827]
- [35] Incze, M.; Redberg, R.F.; Gupta, A. I Have Insomnia-What Should I Do? *JAMA Intern. Med.*, **2018**, 178(11), 1572.  
[http://dx.doi.org/10.1001/jamainternmed.2018.2626] [PMID: 30208397]
- [36] Patel, D.; Steinberg, J.; Patel, P. Insomnia in the Elderly: A Review. *J. Clin. Sleep Med.*, **2018**, 14(6), 1017-1024.  
[http://dx.doi.org/10.5664/jcsm.7172] [PMID: 29852897]
- [37] Sindi, S.; Kåreholt, I.; Johansson, L.; Skoog, J.; Sjöberg, L.; Wang, H.X.; Johansson, B.; Fratiglioni, L.; Soininen, H.; Solomon, A.; Skoog, I.; Kivipelto, M. Sleep disturbances and dementia risk: A multicenter study. *Alzheimer's Dement.*, **2018**, 14(10), 1235-1242.  
[http://dx.doi.org/10.1016/j.jalz.2018.05.012] [PMID: 30030112]
- [38] Desaulniers, J.; Desjardins, S.; Lapierre, S.; Desgagné, A. Sleep Environment and Insomnia in Elderly Persons Living at Home. *J. Aging Res.*, **2018**, 2018, 8053696.  
[http://dx.doi.org/10.1155/2018/8053696] [PMID: 30363712]
- [39] Abad, V.C.; Guilleminault, C. Insomnia in Elderly Patients: Recommendations for Pharmacological Management. *Drugs Aging*, **2018**, 35(9), 791-817.  
[http://dx.doi.org/10.1007/s40266-018-0569-8] [PMID: 30058034]
- [40] Brandão, G.S.; Camelier, F.W.R.; Sampaio, A.A.C.; Brandão, G.S.; Silva, A.S.; Gomes, G.S.B.F.; Donner, C.F.; Oliveira, L.V.F.; Camelier, A.A. Association of sleep quality with excessive daytime somnolence and quality of life of elderlies of community. *Multidiscip. Respir. Med.*, **2018**, 13, 8.  
[http://dx.doi.org/10.1186/s40248-018-0120-0] [PMID: 29568522]
- [41] Garbarino, S.; Scoditti, E.; Lanteri, P.; Conte, L.; Magnavita, N.; Toraldo, D.M. Obstructive Sleep Apnea With or Without Excessive Daytime Sleepiness: Clinical and Experimental Data-Driven Phenotyping. *Front. Neurol.*, **2018**, 9, 505.  
[http://dx.doi.org/10.3389/fneur.2018.00505] [PMID: 29997573]
- [42] Kim, J.W.; Kim, T.; Shin, J.; Choe, G.; Lim, H.J.; Rhee, C.S.; Lee, K.; Cho, S.W. Prediction of Obstructive Sleep Apnea Based on Respiratory Sounds Recorded Between Sleep Onset and Sleep Offset. *Clin. Exp. Otorhinolaryngol.*, **2019**, 12(1), 72-78.  
[http://dx.doi.org/10.21053/ceo.2018.00388] [PMID: 30189718]
- [43] Carneiro-Barrera, A.; Díaz-Román, A.; Guillén-Riquelme, A.; Bucla-Casal, G. Weight loss and lifestyle interventions for obstructive sleep apnoea in adults: Systematic review and meta-analysis. *Obes. Rev.*, **2019**, 20(5), 750-762.  
[http://dx.doi.org/10.1111/obr.12824] [PMID: 30609450]
- [44] Zalai, D.; Bingeliene, A.; Shapiro, C. Sleepiness in the Elderly. *Sleep Med. Clin.*, **2017**, 12(3), 429-441.  
[http://dx.doi.org/10.1016/j.jsmc.2017.03.015] [PMID: 28778240]
- [45] Junho, B.T.; Kummer, A.; Cardoso, F.; Teixeira, A.L.; Rocha, N.P. Clinical Predictors of Excessive Daytime Sleepiness in Patients with Parkinson's Disease. *J. Clin. Neurol.*, **2018**, 14(4), 530-536.  
[http://dx.doi.org/10.3988/jcn.2018.14.4.530] [PMID: 30198233]
- [46] Maugeri, A.; Medina-Inojosa, J.R.; Kunzova, S.; Agodi, A.; Barchitta, M.; Sochor, O.; Lopez-Jimenez, F.; Geda, Y.E.; Vinciguerra, M. Sleep Duration and Excessive Daytime Sleepiness Are Associated with Obesity Independent of Diet and Physical Activity. *Nutrients*, **2018**, 10(9), E1219.  
[http://dx.doi.org/10.3390/nu10091219] [PMID: 30177634]
- [47] Hombali, A.; Seow, E.; Yuan, Q.; Chang, S.H.S.; Satghare, P.; Kumar, S.; Verma, S.K.; Mok, Y.M.; Chong, S.A.; Subramaniam, M. Prevalence and correlates of sleep disorder symptoms in psychiatric disorders. *Psychiatry Res.*, **2018**, S0165-1781(18)30268-3.  
[http://dx.doi.org/10.1016/j.psychres.2018.07.009] [PMID: 30072039]
- [48] K Pavlova M, Latreille V. Sleep Disorders.. *Am J Med*, **2018**. pii: S0002-9343(18)30944-6
- [49] Gulia, K.K.; Kumar, V.M. Sleep disorders in the elderly: a growing challenge. *Psychogeriatrics*, **2018**, 18(3), 155-165.  
[http://dx.doi.org/10.1111/psyg.12319] [PMID: 29878472]
- [50] Iranzo, A. Parasomnias and Sleep-Related Movement Disorders in Older Adults. *Sleep Med. Clin.*, **2018**, 13(1), 51-61.  
[http://dx.doi.org/10.1016/j.jsmc.2017.09.005] [PMID: 29412983]
- [51] Miner, B.; Kryger, M.H. Sleep in the Aging Population. *Sleep Med. Clin.*, **2017**, 12(1), 31-38.  
[http://dx.doi.org/10.1016/j.jsmc.2016.10.008] [PMID: 28159095]
- [52] Grandner, M.A.; Winkelman, J.W. Nocturnal leg cramps: Prevalence and associations with demographics, sleep disturbance symptoms, medical conditions, and cardiometabolic risk factors. *PLoS One*, **2017**, 12(6), e0178465.  
[http://dx.doi.org/10.1371/journal.pone.0178465] [PMID: 28586374]
- [53] Yaremchuk, K. Sleep Disorders in the Elderly. *Clin. Geriatr. Med.*, **2018**, 34(2), 205-216.  
[http://dx.doi.org/10.1016/j.cger.2018.01.008] [PMID: 29661333]
- [54] Wang, C.; Wang, Q.; Ji, B.; Pan, Y.; Xu, C.; Cheng, B.; Bai, B.; Chen, J. The Orexin/Receptor System: Molecular Mechanism and Therapeutic Potential for Neurological Diseases. *Front. Mol. Neurosci.*, **2018**, 11, 220.  
[http://dx.doi.org/10.3389/fnmol.2018.00220] [PMID: 30002617]
- [55] Bonvalet, M.; Ollila, H.M.; Ambati, A.; Mignot, E. Autoimmunity in narcolepsy. *Curr. Opin. Pulm. Med.*, **2017**, 23(6), 522-529.  
[http://dx.doi.org/10.1097/MCP.0000000000000426] [PMID: 28991006]
- [56] Takenoshita, S.; Sakai, N.; Chiba, Y.; Matsumura, M.; Yamaguchi, M.; Nishino, S. An overview of hypocretin based therapy in narcolepsy. *Expert Opin. Investig. Drugs*, **2018**, 27(4), 389-406.  
[http://dx.doi.org/10.1080/13543784.2018.1459561] [PMID: 29623725]
- [57] Fronczek, R.; van Geest, S.; Frölich, M.; Overeem, S.; Roelandse, F.W.; Lammers, G.J.; Swaab, D.F. Hypocretin (orexin) loss in Alzheimer's disease. *Neurobiol. Aging*, **2012**, 33(8), 1642-1650.  
[http://dx.doi.org/10.1016/j.neurobiolaging.2011.03.014] [PMID: 21546124]
- [58] Roh, J.H.; Jiang, H.; Finn, M.B.; Stewart, F.R.; Mahan, T.E.; Cirrito, J.R.; Heda, A.; Snider, B.J.; Li, M.; Yanagisawa, M.; de Lecea, L.; Holtzman, D.M. Potential role of orexin and sleep modulation in the pathogenesis of Alzheimer's disease. *J. Exp. Med.*, **2014**, 211(13), 2487-2496.  
[http://dx.doi.org/10.1084/jem.20141788] [PMID: 25422493]
- [59] Nixon, J.P.; Mavanji, V.; Butterick, T.A.; Billington, C.J.; Kotz, C.M.; Teske, J.A. Sleep disorders, obesity, and aging: the role of orexin. *Ageing Res. Rev.*, **2015**, 20, 63-73.  
[http://dx.doi.org/10.1016/j.arr.2014.11.001] [PMID: 25462194]
- [60] Kovalská, P.; Kemlink, D.; Nevšimálová, S.; Maurovich Horvat, E.; Jarolímová, E.; Topinková, E.; Šonka, K. Narcolepsy with cataplexy in patients aged over 60 years: a case-control study. *Sleep Med.*, **2016**, 26, 79-84.  
[http://dx.doi.org/10.1016/j.sleep.2016.05.011] [PMID: 27665501]
- [61] Gabelle, A.; Jaussent, I.; Hirtz, C.; Vialaret, J.; Navucet, S.; Grasselli, C.; Robert, P.; Lehmann, S.; Dauvilliers, Y. Cerebrospinal fluid levels of orexin-A and histamine, and sleep profile within the Alzheimer process. *Neurobiol. Aging*, **2017**, 53, 59-66.  
[http://dx.doi.org/10.1016/j.neurobiolaging.2017.01.011] [PMID: 28235679]
- [62] Suzuki, K.; Miyamoto, M.; Hirata, K. Sleep disorders in the elderly: Diagnosis and management. *J Gen Fam Med*, **2017**, 18(2), 61-71.  
[http://dx.doi.org/10.1002/jgf2.27] [PMID: 29263993]
- [63] Salas-Crisostomo, M.; Tortorolo, P.; Veras, A.B.; Rocha, N.B.; Machado, S.; Murillo-Rodríguez, E. Therapeutic approaches for the management of sleep disorders in geriatric population. *Curr. Med. Chem.*, **2018**.  
[http://dx.doi.org/10.2174/0929867325666180904113115] [PMID: 30182852]
- [64] Augustin, S.M.; Lovinger, D.M. Functional Relevance of Endocannabinoid-Dependent Synaptic Plasticity in the Central Nervous System. *ACS Chem. Neurosci.*, **2018**, 9(9), 2146-2161.  
[http://dx.doi.org/10.1021/acschemneuro.7b00508] [PMID: 29400439]
- [65] Aymerich, M.S.; Aso, E.; Abellanas, M.A.; Tolón, R.M.; Ramos, J.A.; Ferrer, I.; Romero, J.; Fernández-Ruiz, J. Cannabinoid pharmacology/therapeutics in chronic degenerative disorders affecting the central nervous system. *Biochem. Pharmacol.*, **2018**, 157, 67-84.  
[http://dx.doi.org/10.1016/j.bcp.2018.08.016] [PMID: 30121249]
- [66] Ibarra-Lecue, I.; Pilar-Cuellar, F.; Muguza, C.; Florensa-Zanuy, E.; Díaz, A.; Urigüen, L.; Castro, E.; Pazos, A.; Callado, L.F. The endocannabinoid system in mental disorders: Evidence from human brain studies. *Biochem. Pharmacol.*, **2018**, 157, 97-107.  
[http://dx.doi.org/10.1016/j.bcp.2018.07.009] [PMID: 30026022]



- [67] Jordan, C.J.; Xi, Z.X. Progress in Brain Cannabinoid CB2 Receptor Research: From Genes to Behavior *Neurosci Biobehav Rev*, **2019**, pii: S0149-7634(18)30829-7
- [68] Hu, S.S.; Mackie, K. Distribution of the Endocannabinoid System in the Central Nervous System. *Handb. Exp. Pharmacol.*, **2015**, *231*, 59-93. [http://dx.doi.org/10.1007/978-3-319-20825-1\_3] [PMID: 26408158]
- [69] Kendall, D.A.; Yudowski, G.A. Cannabinoid Receptors in the Central Nervous System: Their Signaling and Roles in Disease. *Front. Cell. Neurosci.*, **2017**, *10*, 294. [http://dx.doi.org/10.3389/fncel.2016.00294] [PMID: 28101004]
- [70] Navarro, G.; Morales, P.; Rodríguez-Cueto, C.; Fernández-Ruiz, J.; Jagerovic, N.; Franco, R. Targeting Cannabinoid CB2 Receptors in the Central Nervous System. Medicinal Chemistry Approaches with Focus on Neurodegenerative Disorders. *Front. Neurosci.*, **2016**, *10*, 406. [http://dx.doi.org/10.3389/fnins.2016.00406] [PMID: 27679556]
- [71] Bloomfield, M.P.; Hindocha, C.; Green, S.F.; Wall, M.B.; Lees, R.; Petrilli, K.; Costello, H.; Ogunbiyi, M.O.; Bossong, M.G.; Freeman, T.P. The neuropsychopharmacology of cannabis: A review of human imaging studies. *Pharmacol Ther.*, **2018**, pii: S0163-7258(18)30190-6
- [72] Maccarrone, M.; Bab, I.; Biró, T.; Cabral, G.A.; Dey, S.K.; Di Marzo, V.; Konje, J.C.; Kunos, G.; Mechoulam, R.; Pacher, P.; Sharkey, K.A.; Zimmer, A. Endocannabinoid signaling at the periphery: 50 years after THC. *Trends Pharmacol. Sci.*, **2015**, *36*(5), 277-296. [http://dx.doi.org/10.1016/j.tips.2015.02.008] [PMID: 25796370]
- [73] Deshmukh, R.R.; Sharma, P.L. Stimulation of accumbens shell cannabinoid CB1 receptors by noladin ether, a putative endocannabinoid, modulates food intake and dietary selection in rats. *Pharmacol. Res.*, **2012**, *66*(3), 276-282. [http://dx.doi.org/10.1016/j.phrs.2012.06.004] [PMID: 22728691]
- [74] Pertwee, R.G. Endocannabinoids and Their Pharmacological Actions. *Handb. Exp. Pharmacol.*, **2015**, *231*, 1-37. [http://dx.doi.org/10.1007/978-3-319-20825-1\_1] [PMID: 26408156]
- [75] Redmond, W.J.; Cawston, E.E.; Grimsey, N.L.; Stuart, J.; Edington, A.R.; Glass, M.; Connor, M. Identification of N-arachidonoyl dopamine as a highly biased ligand at cannabinoid CB1 receptors. *Br. J. Pharmacol.*, **2016**, *173*(1), 115-127. [http://dx.doi.org/10.1111/bph.13341] [PMID: 26398720]
- [76] Lawton, S.K.; Xu, F.; Tran, A.; Wong, E.; Prakash, A.; Schumacher, M.; Hellman, J.; Wilhelmsen, K. N-Arachidonoyl Dopamine Modulates Acute Systemic Inflammation via Nonhematopoietic TRPV1. *J. Immunol.*, **2017**, *199*(4), 1465-1475. [http://dx.doi.org/10.4049/jimmunol.1602151] [PMID: 28701511]
- [77] Pandey, P.; Chaurasiya, N.D.; Tekwani, B.L.; Doerksen, R.J. Interactions of endocannabinoid virodhamine and related analogs with human monoamine oxidase-A and -B. *Biochem. Pharmacol.*, **2018**, *155*, 82-91. [http://dx.doi.org/10.1016/j.bcp.2018.06.024] [PMID: 29958841]
- [78] Chanda, D.; Neumann, D.; Glatz, J.F.C. The endocannabinoid system: Overview of an emerging multi-faceted therapeutic target. *Prostaglandins Leukot. Essent. Fatty Acids*, **2019**, *140*, 51-56. [http://dx.doi.org/10.1016/j.plefa.2018.11.016] [PMID: 30553404]
- [79] Maccarrone, M. Metabolism of the Endocannabinoid Anandamide: Open Questions after 25 Years. *Front. Mol. Neurosci.*, **2017**, *10*, 166. [http://dx.doi.org/10.3389/fnmol.2017.00166] [PMID: 28611591]
- [80] Nazıroğlu, M.; Taner, A.N.; Balbay, E.; Çiğ, B. Inhibitions of anandamide transport and FAAH synthesis decrease apoptosis and oxidative stress through inhibition of TRPV1 channel in an *in vitro* seizure model. *Mol. Cell. Biochem.*, **2019**, *453*(1-2), 143-155. [http://dx.doi.org/10.1007/s11010-018-3439-0] [PMID: 30159798]
- [81] Storozhuk, M.V.; Zholos, A.V. TRP Channels as Novel Targets for Endogenous Ligands: Focus on Endocannabinoids and Nociceptive Signalling. *Curr. Neuropharmacol.*, **2018**, *16*(2), 137-150. [http://dx.doi.org/10.2174/1570159X15666170424120802] [PMID: 28440188]
- [82] Toczek, M.; Malinowska, B. Enhanced endocannabinoid tone as a potential target of pharmacotherapy. *Life Sci.*, **2018**, *204*, 20-45. [http://dx.doi.org/10.1016/j.lfs.2018.04.054] [PMID: 29729263]
- [83] Patel, S.; Hill, M.N.; Cheer, J.F.; Wotjak, C.T.; Holmes, A. The endocannabinoid system as a target for novel anxiolytic drugs. *Neurosci Biobehav Rev*, **2017**, *76*(Pt A), 56-66. [http://dx.doi.org/10.1016/j.neubiorev.2016.12.033]
- [84] Burstein, S.H. N-Acyl Amino Acids (Elmiric Acids): Endogenous Signaling Molecules with Therapeutic Potential. *Mol. Pharmacol.*, **2018**, *93*(3), 228-238. [http://dx.doi.org/10.1124/mol.117.110841] [PMID: 29138268]
- [85] Morena, M.; Aukema, R.J.; Leilt, K.D.; Rashid, A.J.; Vecchiarelli, H.A.; Josselyn, S.A.; Hill, M.N. Upregulation of Anandamide Hydrolysis in the Basolateral Complex of Amygdala Reduces Fear Memory Expression and Indices of Stress and Anxiety. *J. Neurosci.*, **2018**, 2251-22518. [PMID: 30573646]
- [86] Jumpertz, R.; Wiesner, T.; Blüher, M.; Engeli, S.; Bátkai, S.; Wirtz, H.; Bosse-Henck, A.; Stumvoll, M. Circulating endocannabinoids and N-acyl-ethanolamides in patients with sleep apnea--specific role of oleylethanolamide. *Exp. Clin. Endocrinol. Diabetes*, **2010**, *118*(9), 591-595. [http://dx.doi.org/10.1055/s-0030-1253344] [PMID: 20429051]
- [87] Murillo-Rodríguez, E.; Poot-Ake, A.; Arias-Carrion, O.; Pacheco-Pantoja, E.; Fuente-Ortega, A. de L.; Arankowsky-Sandoval, G. The emerging role of the endocannabinoid system in the sleep-wake cycle modulation. *Cent. Nerv. Syst. Agents Med. Chem.*, **2011**, *11*(3), 189-196. [http://dx.doi.org/10.2174/187152411798047780] [PMID: 21919868]
- [88] Engeli, S.; Blüher, M.; Jumpertz, R.; Wiesner, T.; Wirtz, H.; Bosse-Henck, A.; Stumvoll, M.; Bátkai, S.; Pacher, P.; Harvey-White, J.; Kunos, G.; Jordan, J. Circulating anandamide and blood pressure in patients with obstructive sleep apnea. *J. Hypertens.*, **2012**, *30*(12), 2345-2351. [http://dx.doi.org/10.1097/HJH.0b013e3283591595] [PMID: 23032139]
- [89] Murillo-Rodríguez, E.; Palomero-Rivero, M.; Millán-Aldaco, D.; Di Marzo, V. The administration of endocannabinoid uptake inhibitors OMDM-2 or VDM-11 promotes sleep and decreases extracellular levels of dopamine in rats. *Physiol. Behav.*, **2013**, *109*, 88-95. [http://dx.doi.org/10.1016/j.physbeh.2012.11.007] [PMID: 23238438]
- [90] Murillo-Rodríguez, E.; Machado, S.; Rocha, N.B.; Budde, H.; Yuan, T.F.; Arias-Carrion, O. Revealing the role of the endocannabinoid system modulators, SR141716A, URB597 and VDM-11, in sleep homeostasis. *Neuroscience*, **2016**, *339*, 433-449. [http://dx.doi.org/10.1016/j.neuroscience.2016.10.011] [PMID: 27746343]
- [91] Hanlon, E.C.; Tasali, E.; Leproult, R.; Stuhr, K.L.; Doncheck, E.; de Wit, H.; Hillard, C.J.; Van Cauter, E. Sleep Restriction Enhances the Daily Rhythm of Circulating Levels of Endocannabinoid 2-Arachidonoylglycerol. *Sleep (Basel)*, **2016**, *39*(3), 653-664. [http://dx.doi.org/10.5665/sleep.5546] [PMID: 26612385]
- [92] Prospero-García, O.; Amancio-Belmont, O.; Becerril Meléndez, A.L.; Ruiz-Contreras, A.E.; Méndez-Díaz, M. Endocannabinoids and sleep. *Neurosci. Biobehav. Rev.*, **2016**, *71*, 671-679. [http://dx.doi.org/10.1016/j.neubiorev.2016.10.005] [PMID: 27756691]
- [93] Soni, N.; Prabhala, B.K.; Mehta, V.; Mirza, O.; Kohlmeier, K.A. Anandamide and 2-AG are endogenously present within the laterodorsal tegmental nucleus: Functional implications for a role of eCBs in arousal. *Brain Res.*, **2017**, *1665*, 74-79. [http://dx.doi.org/10.1016/j.brainres.2017.04.003] [PMID: 28404451]
- [94] Murguruza, C.; Lehtonen, M.; Aaltonen, N.; Morentin, B.; Meana, J.J.; Callado, L.F. Quantification of endocannabinoids in postmortem brain of schizophrenic subjects. *Schizophr. Res.*, **2013**, *148*(1-3), 145-150. [http://dx.doi.org/10.1016/j.schres.2013.06.013] [PMID: 23800614]
- [95] Rodríguez-Cueto, C.; Hernández-Gálvez, M.; Hillard, C.J.; Maciel, P.; Valdeolivas, S.; Ramos, J.A.; Gómez-Ruiz, M.; Fernández-Ruiz, J. Altered striatal endocannabinoid signaling in a transgenic mouse model of spinocerebellar ataxia type-3. *PLoS One*, **2017**, *12*(4), e0176521. [http://dx.doi.org/10.1371/journal.pone.0176521] [PMID: 28448548]

- [96] Micale, V.; Drago, F. Endocannabinoid system, stress and HPA axis. *Eur. J. Pharmacol.*, **2018**, *834*, 230-239. [http://dx.doi.org/10.1016/j.ejphar.2018.07.039] [PMID: 30036537]
- [97] Aso, E.; Andrés-Benito, P.; Ferrer, I. Genetic deletion of CB<sub>1</sub> cannabinoid receptors exacerbates the Alzheimer-like symptoms in a transgenic animal model. *Biochem. Pharmacol.*, **2018**, *157*, 210-216. [http://dx.doi.org/10.1016/j.bcp.2018.08.007] [PMID: 30096288]
- [98] Choi, S.H.; Arai, A.L.; Mou, Y.; Kang, B.; Yen, C.C.; Hallenbeck, J.; Silva, A.C. Neuroprotective Effects of MAGL (Monoacylglycerol Lipase) Inhibitors in Experimental Ischemic Stroke. *Stroke*, **2018**, *49*(3), 718-726. [http://dx.doi.org/10.1161/STROKEAHA.117.019664] [PMID: 29440474]
- [99] Ilyasov, A.A.; Milligan, C.E.; Pharr, E.P.; Howlett, A.C. The Endocannabinoid System and Oligodendrocytes in Health and Disease. *Front. Neurosci.*, **2018**, *12*, 733. [http://dx.doi.org/10.3389/fnins.2018.00733] [PMID: 30416422]
- [100] Iannotti, F.A.; Pagano, E.; Guardiola, O.; Adinolfi, S.; Saccone, V.; Consalvi, S.; Piscitelli, F.; Gazzero, E.; Busetto, G.; Carrella, D.; Capasso, R.; Puri, P.L.; Minchiotti, G.; Di Marzo, V. Genetic and pharmacological regulation of the endocannabinoid CB<sub>1</sub> receptor in Duchenne muscular dystrophy. *Nat. Commun.*, **2018**, *9*(1), 3950. [http://dx.doi.org/10.1038/s41467-018-06267-1] [PMID: 30262909]
- [101] Marchioni, C.; de Souza, I.D.; Acquaro, V.R.; de Souza Crippa, J.A.; Tumas, V.; Queiroz, M.E.C. Recent advances in LC-MS/MS methods to determine endocannabinoids in biological samples: Application in neurodegenerative diseases. *Anal. Chim. Acta*, **2018**, *1044*, 12-28. [http://dx.doi.org/10.1016/j.aca.2018.06.016] [PMID: 30442393]
- [102] Mounsey, R.B.; Mustafa, S.; Robinson, L.; Ross, R.A.; Riedel, G.; Pertwee, R.G.; Teismann, P. Increasing levels of the endocannabinoid 2-AG is neuroprotective in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine mouse model of Parkinson's disease. *Exp. Neurol.*, **2015**, *273*, 36-44. [http://dx.doi.org/10.1016/j.expneurol.2015.07.024] [PMID: 26244281]
- [103] Basavarajappa, B.S.; Shivakumar, M.; Joshi, V.; Subbanna, S. Endocannabinoid system in neurodegenerative disorders. *J. Neurochem.*, **2017**, *142*(5), 624-648. [http://dx.doi.org/10.1111/jnc.14098] [PMID: 28608560]
- [104] D'Addario, C.; Di Francesco, A.; Arosio, B.; Gussago, C.; Dell'Osso, B.; Bari, M.; Galimberti, D.; Scarpini, E.; Altamura, A.C.; Mari, D.; Maccarrone, M. Epigenetic regulation of fatty acid amide hydrolase in Alzheimer disease. *PLoS One*, **2012**, *7*(6), e39186. [http://dx.doi.org/10.1371/journal.pone.0039186] [PMID: 22720070]
- [105] Pascual, A.C.; Martín-Moreno, A.M.; Giusto, N.M.; de Ceballos, M.L.; Pasquaré, S.J. Normal aging in rats and pathological aging in human Alzheimer's disease decrease FAAH activity: modulation by cannabinoid agonists. *Exp. Gerontol.*, **2014**, *60*, 92-99. [http://dx.doi.org/10.1016/j.exger.2014.10.011] [PMID: 25456842]
- [106] Bedse, G.; Romano, A.; Lavecchia, A.M.; Cassano, T.; Gaetani, S. The role of endocannabinoid signaling in the molecular mechanisms of neurodegeneration in Alzheimer's disease. *J. Alzheimers Dis.*, **2015**, *43*(4), 1115-1136. [http://dx.doi.org/10.3233/JAD-141635] [PMID: 25147120]
- [107] Szűcs, E.; Dvoračská, S.; Tömböly, C.; Büki, A.; Kékesi, G.; Horváth, G.; Benyhe, S. Decreased CB receptor binding and cannabinoid signaling in three brain regions of a rat model of schizophrenia. *Neurosci. Lett.*, **2016**, *633*, 87-93. [http://dx.doi.org/10.1016/j.neulet.2016.09.020] [PMID: 27639959]
- [108] Fakhoury, M. Role of the Endocannabinoid System in the Pathophysiology of Schizophrenia. *Mol. Neurobiol.*, **2017**, *54*(1), 768-778. [http://dx.doi.org/10.1007/s12035-016-9697-5] [PMID: 26768595]
- [109] Ibarra-Lecue, I.; Pilar-Cuellar, F.; Muguruza, C.; Florensa-Zanuy, E.; Díaz, Á.; Urigüen, L.; Castro, E.; Pazos, A.; Callado, L.F. The endocannabinoid system in mental disorders: Evidence from human brain studies. *Biochem. Pharmacol.*, **2018**, *157*, 97-107. [http://dx.doi.org/10.1016/j.bcp.2018.07.009] [PMID: 30026022]
- [110] Bioque, M.; García-Bueno, B.; Macdowell, K.S.; Meseguer, A.; Saiz, P.A.; Parellada, M.; Gonzalez-Pinto, A.; Rodriguez-Jimenez, R.; Lobo, A.; Leza, J.C.; Bernardo, M. FLAMM-PEPs study—Centro de Investigación Biomedica en Red de Salud Mental. Peripheral endocannabinoid system dysregulation in first-episode psychosis. *Neuropsychopharmacol.*, **2013**, *38*(13), 2568-2577. [http://dx.doi.org/10.1038/npp.2013.165] [PMID: 23811653]
- [111] Santucci, V.; Storme, J.J.; Soubrié, P.; Le Fur, G. Arousal-enhancing properties of the CB<sub>1</sub> cannabinoid receptor antagonist SR 141716A in rats as assessed by electroencephalographic spectral and sleep-waking cycle analysis. *Life Sci.*, **1996**, *58*(6), PL103-PL110. [http://dx.doi.org/10.1016/0024-3205(95)02319-4] [PMID: 8569415]
- [112] Murillo-Rodríguez, E.; Sánchez-Alavez, M.; Navarro, L.; Martínez-González, D.; Drucker-Colín, R.; Prospéro-García, O. Anandamide modulates sleep and memory in rats. *Brain Res.*, **1998**, *812*(1-2), 270-274. [http://dx.doi.org/10.1016/S0006-8993(98)00969-X] [PMID: 9813364]
- [113] Murillo-Rodríguez, E.; Cabeza, R.; Méndez-Díaz, M.; Navarro, L.; Prospéro-García, O. Anandamide-induced sleep is blocked by SR141716A, a CB<sub>1</sub> receptor antagonist and by U73122, a phospholipase C inhibitor. *Neuroreport*, **2001**, *12*(10), 2131-2136. [http://dx.doi.org/10.1097/00001756-200107200-00018] [PMID: 11447321]
- [114] Murillo-Rodríguez, E.; Vázquez, E.; Millán-Aldaco, D.; Palomero-Rivero, M.; Drucker-Colín, R. Effects of the fatty acid amide hydrolase inhibitor URB597 on the sleep-wake cycle, c-Fos expression and dopamine levels of the rat. *Eur. J. Pharmacol.*, **2007**, *562*(1-2), 82-91. [http://dx.doi.org/10.1016/j.ejphar.2007.01.076] [PMID: 17336288]
- [115] Murillo-Rodríguez, E.; Palomero-Rivero, M.; Millán-Aldaco, D.; Arias-Carrión, O.; Drucker-Colín, R. Administration of URB597, oleoylethanolamide or palmitoylethanolamide increases waking and dopamine in rats. *PLoS One*, **2011**, *6*(7), e20766. [http://dx.doi.org/10.1371/journal.pone.0020766] [PMID: 21779318]
- [116] Tutunchi, H.; Ostadrahimi, A.; Saghaei-Asl, M.; Maleki, V. The effects of oleoylethanolamide, an endogenous PPAR- $\alpha$  agonist, on risk factors for NAFLD: A systematic review. *Obes. Rev.*, **2019**, *20*(7), 1057-1069. [http://dx.doi.org/10.1111/obr.12853] [PMID: 31111657]
- [117] Mijangos-Moreno, S.; Poot-Aké, A.; Guzmán, K.; Arankowsky-Sandoval, G.; Arias-Carrión, O.; Zaldivar-Rae, J.; Sarro-Ramírez, A.; Murillo-Rodríguez, E. Sleep and neurochemical modulation by the nuclear peroxisome proliferator-activated receptor  $\alpha$  (PPAR- $\alpha$ ) in rat. *Neurosci. Res.*, **2016**, *105*, 65-69. [http://dx.doi.org/10.1016/j.neures.2015.09.005] [PMID: 26450400]
- [118] Murillo-Rodríguez, E. The Role of Nuclear Receptor PPAR $\alpha$  in the Sleep-wake Cycle Modulation. A Tentative Approach for Treatment of Sleep Disorders. *Curr. Drug Deliv.*, **2017**, *14*(4), 473-482. [http://dx.doi.org/10.2174/1567201814666161109123803] [PMID: 27834148]
- [119] Murillo-Rodríguez, E.; Millán-Aldaco, D.; Di Marzo, V.; Drucker-Colín, R. The anandamide membrane transporter inhibitor, VDM-11, modulates sleep and c-Fos expression in the rat brain. *Neuroscience*, **2008**, *157*(1), 1-11. [http://dx.doi.org/10.1016/j.neuroscience.2008.08.056] [PMID: 18822353]
- [120] Murillo-Rodríguez, E.; Palomero-Rivero, M.; Millán-Aldaco, D.; Di Marzo, V. The administration of endocannabinoid uptake inhibitors OMDM-2 or VDM-11 promotes sleep and decreases extracellular levels of dopamine in rats. *Physiol. Behav.*, **2013**, *109*, 88-95. [http://dx.doi.org/10.1016/j.physbeh.2012.11.007] [PMID: 23238438]
- [121] Murillo-Rodríguez, E.; Machado, S.; Rocha, N.B.; Budde, H.; Yuan, T.F.; Arias-Carrión, O. Revealing the role of the endocannabinoid system modulators, SR141716A, URB597 and VDM-11, in sleep homeostasis. *Neuroscience*, **2016**, *339*, 433-449. [http://dx.doi.org/10.1016/j.neuroscience.2016.10.011] [PMID: 27746343]

- [122] Engeli, S.; Blüher, M.; Jumpertz, R.; Wiesner, T.; Wirtz, H.; Bosse-Henck, A.; Stumvoll, M.; Batkai, S.; Pacher, P.; Harvey-White, J.; Kunos, G.; Jordan, J. Circulating anandamide and blood pressure in patients with obstructive sleep apnea. *J. Hypertens.*, **2012**, *30*(12), 2345-2351. [http://dx.doi.org/10.1097/HJH.0b013e3283591595] [PMID: 23032139]
- [123] Wang, X.; Yu, Q.; Yue, H.; Zhang, J.; Zeng, S.; Cui, F. Circulating Endocannabinoids and Insulin Resistance in Patients with Obstructive Sleep Apnea. *BioMed Res. Int.*, **2016**, *2016*, 9782031. [http://dx.doi.org/10.1155/2016/9782031] [PMID: 26904688]
- [124] Maple, K.E.; McDaniel, K.A.; Shollenbarger, S.G.; Lisdahl, K.M. Dose-dependent cannabis use, depressive symptoms, and FAAH genotype predict sleep quality in emerging adults: a pilot study. *Am. J. Drug Alcohol Abuse*, **2016**, *42*(4), 431-440. [http://dx.doi.org/10.3109/00952990.2016.1141913] [PMID: 27074158]
- [125] Bilkei-Gorzo, A. The endocannabinoid system in normal and pathological brain ageing. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, **2012**, *367*(1607), 3326-3341. [http://dx.doi.org/10.1098/rstb.2011.0388] [PMID: 23108550]
- [126] Piyanova, A.; Lomazzo, E.; Bindila, L.; Lerner, R.; Albayram, O.; Ruhl, T.; Lutz, B.; Zimmer, A.; Bilkei-Gorzo, A. Age-related changes in the endocannabinoid system in the mouse hippocampus. *Mech. Ageing Dev.*, **2015**, *150*, 55-64. [http://dx.doi.org/10.1016/j.mad.2015.08.005] [PMID: 26278494]
- [127] Bonnet, A.E.; Marchalant, Y. Potential Therapeutic Contributions of the Endocannabinoid System towards Aging and Alzheimer's Disease. *Aging Dis.*, **2015**, *6*(5), 400-405. [http://dx.doi.org/10.14336/AD.2015.0617] [PMID: 26425394]
- [128] Vázquez, C.; Tolón, R.M.; Grande, M.T.; Caraza, M.; Moreno, M.; Koester, E.C.; Villacusa, B.; Ruiz-Valdepeñas, L.; Fernández-Sánchez, F.J.; Cravatt, B.F.; Hillard, C.J.; Romero, J. Endocannabinoid regulation of amyloid-induced neuroinflammation. *Neurobiol. Aging*, **2015**, *36*(11), 3008-3019. [http://dx.doi.org/10.1016/j.neurobiolaging.2015.08.003] [PMID: 26362942]
- [129] Sophocleous, A.; Marino, S.; Kabir, D.; Ralston, S.H.; Idris, A.I. Combined deficiency of the Cnr1 and Cnr2 receptors protects against age-related bone loss by osteoclast inhibition. *Aging Cell*, **2017**, *16*(5), 1051-1061. [http://dx.doi.org/10.1111/acer.12638] [PMID: 28752643]
- [130] Cassano, T.; Calcagnini, S.; Pace, L.; De Marco, F.; Romano, A.; Gaetani, S. Cannabinoid Receptor 2 Signaling in Neurodegenerative Disorders: From Pathogenesis to a Promising Therapeutic Target. *Front. Neurosci.*, **2017**, *11*, 30. [http://dx.doi.org/10.3389/fnins.2017.00030] [PMID: 28210207]
- [131] Bilkei-Gorzo, A.; Albayram, O.; Draffehn, A.; Michel, K.; Piyanova, A.; Oppenheimer, H.; Dvir-Ginzberg, M.; Rácz, I.; Ulas, T.; Imbeault, S.; Bab, I.; Schultze, J.L.; Zimmer, A. A chronic low dose of  $\Delta^9$ -tetrahydrocannabinol (THC) restores cognitive function in old mice. *Nat. Med.*, **2017**, *23*(6), 782-787. [http://dx.doi.org/10.1038/nm.4311] [PMID: 28481360]
- [132] Canseco-Alba, A.; Schanz, N.; Sanabria, B.; Zhao, J.; Lin, Z.; Liu, Q.R.; Onaivi, E.S. Behavioral effects of psychostimulants in mutant mice with cell-type specific deletion of CB2 cannabinoid receptors in dopamine neurons. *Behav. Brain Res.*, **2019**, *360*, 286-297. [http://dx.doi.org/10.1016/j.bbr.2018.11.043] [PMID: 30508607]
- [133] Berrendero, F.; Romero, J.; García-Gil, L.; Suarez, I.; De la Cruz, P.; Ramos, J.A.; Fernández-Ruiz, J.J. Changes in cannabinoid receptor binding and mRNA levels in several brain regions of aged rats. *Biochim. Biophys. Acta*, **1998**, *1407*(3), 205-214. [http://dx.doi.org/10.1016/S0925-4439(98)00042-8] [PMID: 9748581]
- [134] Valenti, M.; Viganò, D.; Casico, M.G.; Rubino, T.; Steardo, L.; Parolaro, D.; Di Marzo, V. Differential diurnal variations of anandamide and 2-arachidonoyl-glycerol levels in rat brain. *Cell. Mol. Life Sci.*, **2004**, *61*(7-8), 945-950. [http://dx.doi.org/10.1007/s00018-003-3453-5] [PMID: 15095014]
- [135] Murillo-Rodríguez, E.; Désarnaud, F.; Prospéro-García, O. Diurnal variation of arachidonoyl ethanolamine, palmitoylethanolamine and oleoylethanolamine in the brain of the rat. *Life Sci.*, **2006**, *79*(1), 30-37. [http://dx.doi.org/10.1016/j.lfs.2005.12.028] [PMID: 16434061]
- [136] Maroof, N.; Ravipati, S.; Pardon, M.C.; Barrett, D.A.; Kendall, D.A. Reductions in endocannabinoid levels and enhanced coupling of cannabinoid receptors in the striatum are accompanied by cognitive impairments in the AβPPswe/PS1ΔE9 mouse model of Alzheimer's disease. *J. Alzheimers Dis.*, **2014**, *42*(1), 227-245. [http://dx.doi.org/10.3233/JAD-131961] [PMID: 24844690]
- [137] Schaich, C.L.; Shaltout, H.A.; Grabenauer, M.; Thomas, B.F.; Gallagher, P.E.; Howlett, A.C.; Diz, D.I. Alterations in the Medullary Endocannabinoid System Contribute to Age-related Impairment of Baroreflex Sensitivity. *J. Cardiovasc. Pharmacol.*, **2015**, *65*(5), 473-479. [http://dx.doi.org/10.1097/FJC.0000000000000216] [PMID: 25636077]
- [138] Tuo, W.; Leleu-Chavain, N.; Spencer, J.; Sansook, S.; Millet, R.; Chavatte, P. Therapeutic Potential of Fatty Acid Amide Hydrolase, Monoacylglycerol Lipase, and N-Acylethanolamine Acid Amidase Inhibitors. *J. Med. Chem.*, **2017**, *60*(1), 4-46. [http://dx.doi.org/10.1021/acs.jmedchem.6b00538] [PMID: 27766867]
- [139] Pascual, A.C.; Martín-Moreno, A.M.; Giusto, N.M.; de Ceballos, M.L.; Pasquaré, S.J. Normal aging in rats and pathological aging in human Alzheimer's disease decrease FAAH activity: modulation by cannabinoid agonists. *Exp. Gerontol.*, **2014**, *60*, 92-99. [http://dx.doi.org/10.1016/j.exger.2014.10.011] [PMID: 25456842]
- [140] Pascual, A.C.; Gaveglia, V.L.; Giusto, N.M.; Pasquaré, S.J. Aging modifies the enzymatic activities involved in 2-arachidonoylglycerol metabolism. *Biofactors*, **2013**, *39*(2), 209-220. [http://dx.doi.org/10.1002/biof.1055] [PMID: 23281018]
- [141] Pascual, A.C.; Gaveglia, V.L.; Giusto, N.M.; Pasquaré, S.J. Cannabinoid receptor-dependent metabolism of 2-arachidonoylglycerol during aging. *Exp. Gerontol.*, **2014**, *55*, 134-142. [http://dx.doi.org/10.1016/j.exger.2014.04.008] [PMID: 24768821]
- [142] Maccarrone, M.; Attinà, M.; Bari, M.; Carboni, A.; Ledent, C.; Finazzi-Agrò, A. Anandamide degradation and N-acylethanolamines level in wild-type and CB1 cannabinoid receptor knockout mice of different ages. *J. Neurochem.*, **2001**, *78*(2), 339-348. [http://dx.doi.org/10.1046/j.1471-4159.2001.00413.x] [PMID: 11461969]
- [143] Piyanova, A.; Lomazzo, E.; Bindila, L.; Lerner, R.; Albayram, O.; Ruhl, T.; Lutz, B.; Zimmer, A.; Bilkei-Gorzo, A. Age-related changes in the endocannabinoid system in the mouse hippocampus. *Mech. Ageing Dev.*, **2015**, *150*, 55-64. [http://dx.doi.org/10.1016/j.mad.2015.08.005] [PMID: 26278494]
- [144] Lu, H.C.; Mackie, K. An Introduction to the Endogenous Cannabinoid System. *Biol. Psychiatry*, **2016**, *79*(7), 516-525. [http://dx.doi.org/10.1016/j.biopsych.2015.07.028] [PMID: 26698193]
- [145] Boggs, D.L.; Nguyen, J.D.; Morgenson, D.; Taffe, M.A.; Ranganathan, M. Clinical and Preclinical Evidence for Functional Interactions of Cannabidiol and  $\Delta^9$ -Tetrahydrocannabinol. *Neuropsychopharmacology*, **2018**, *43*(1), 142-154. [http://dx.doi.org/10.1038/npp.2017.209] [PMID: 28875990]
- [146] Russo, E.B. Cannabis Therapeutics and the Future of Neurology. *Front. Integr. Neurosci.*, **2018**, *12*, 51. [http://dx.doi.org/10.3389/fnint.2018.00051] [PMID: 30405366]
- [147] Schrot, R.J.; Hubbard, J.R. Cannabinoids: Medical implications. *Ann. Med.*, **2016**, *48*(3), 128-141. [http://dx.doi.org/10.3109/07853890.2016.1145794] [PMID: 26912385]
- [148] World Health Organization 2018. CANNABIDIOL (CBD) Critical Review Report.. <https://www.who.int/medicines/access/controlled-substances/CannabidiolCriticalReview.pdf>
- [149] Ibeas Bih, C.; Chen, T.; Nunn, A.V.; Bazetou, M.; Dallas, M.; Whalley, B.J. Molecular Targets of Cannabidiol in Neurological Disorders. *Neurotherapeutics*, **2015**, *12*(4), 699-730. [http://dx.doi.org/10.1007/s13311-015-0377-3] [PMID: 26264914]
- [150] National Academies of Sciences, Engineering, and Medicine. 2017. The Health Effects of Cannabis and Cannabinoids: The Current State of Evidence and Recommendations for Research. Washington, DC: The National Academies Press. [http://dx.doi.org/10.17226/24625]
- [151] Chagas, M.H.; Eckeli, A.L.; Zuardi, A.W.; Pena-Pereira, M.A.; Sobreira-Neto, M.A.; Sobreira, E.T.; Camilo, M.R.; Bergamaschi, M.M.; Schenck, C.H.; Hallak, J.E.; Tumas, V.; Crippa, J.A. Cannabidiol can improve complex sleep-related behaviours associated

- with rapid eye movement sleep behaviour disorder in Parkinson's disease patients: a case series. *J. Clin. Pharm. Ther.*, **2014**, *39*(5), 564-566. [http://dx.doi.org/10.1111/jcpt.12179] [PMID: 24845114]
- [152] Shannon, S.; Opila-Lehman, J. Effectiveness of Cannabidiol Oil for Pediatric Anxiety and Insomnia as Part of Posttraumatic Stress Disorder: A Case Report. *Perm. J.*, **2016**, *20*(4), 16-005. [http://dx.doi.org/10.7812/TPP/16-005] [PMID: 27768570]
- [153] Shannon, S.; Lewis, N.; Lee, H.; Hughes, S. Cannabidiol in Anxiety and Sleep: A Large Case Series. *Perm. J.*, **2019**, *23*, 18-041. [http://dx.doi.org/10.7812/TPP/18-041] [PMID: 30624194]
- [154] Pertwee, R.G. The diverse CB<sub>1</sub> and CB<sub>2</sub> receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br. J. Pharmacol.*, **2008**, *153*(2), 199-215. [http://dx.doi.org/10.1038/sj.bjp.0707442] [PMID: 17828291]
- [155] Morales, P.; Goya, P.; Jagerovic, N.; Hernandez-Folgado, L. Allosteric Modulators of the CB<sub>1</sub> Cannabinoid Receptor: A Structural Update Review. *Cannabis Cannabinoid Res.*, **2016**, *1*(1), 22-30. [http://dx.doi.org/10.1089/can.2015.0005] [PMID: 28861476]
- [156] Navarro, G.; Reyes-Resina, I.; Rivas-Santisteban, R.; Sánchez de Medina, V.; Morales, P.; Casano, S.; Ferreira-Vera, C.; Lillo, A.; Aguinaga, D.; Jagerovic, N.; Nadal, X.; Franco, R. Cannabidiol skews biased agonism at cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptors with smaller effect in CB<sub>1</sub>-CB<sub>2</sub> heteroreceptor complexes. *Biochem. Pharmacol.*, **2018**, *157*, 148-158. [http://dx.doi.org/10.1016/j.bcp.2018.08.046] [PMID: 30194918]
- [157] Tham, M.; Yilmaz, O.; Alaverdashvili, M.; Kelly, M.E.M.; Denovan-Wright, E.M.; Laprairie, R.B. Allosteric and orthosteric pharmacology of cannabidiol and cannabidiol-dimethylheptyl at the type 1 and type 2 cannabinoid receptors. *Br. J. Pharmacol.*, **2019**, *176*(10), 1455-1469. [http://dx.doi.org/10.1111/bph.14440] [PMID: 29981240]
- [158] Fogaça, M.V.; Campos, A.C.; Coelho, L.D.; Duman, R.S.; Guimarães, F.S. The anxiolytic effects of cannabidiol in chronically stressed mice are mediated by the endocannabinoid system: Role of neurogenesis and dendritic remodeling. *Neuropharmacology*, **2018**, *135*, 22-33. [http://dx.doi.org/10.1016/j.neuropharm.2018.03.001] [PMID: 29510186]
- [159] Mijangos-Moreno, S.; Poot-Aké, A.; Arankowsky-Sandoval, G.; Murillo-Rodríguez, E. Intrahypothalamic injection of cannabidiol increases the extracellular levels of adenosine in nucleus accumbens in rats. *Neurosci. Res.*, **2014**, *84*, 60-63. [http://dx.doi.org/10.1016/j.neures.2014.04.006] [PMID: 24800644]
- [160] Murillo-Rodríguez, E.; Di Marzo, V.; Machado, S.; Rocha, N.B.; Veras, A.B.; Neto, G.A.M.; Budde, H.; Arias-Carrión, O.; Arankowsky-Sandoval, G. Role of N-Arachidonoyl-Serotonin (AA-5-HT) in Sleep-Wake Cycle Architecture, Sleep Homeostasis, and Neurotransmitters Regulation. *Front. Mol. Neurosci.*, **2017**, *10*, 152. [http://dx.doi.org/10.3389/fnmol.2017.00152] [PMID: 28611585]
- [161] Murillo-Rodríguez, E.; Arankowsky-Sandoval, G.; Rocha, N.B.; Peniche-Amante, R.; Veras, A.B.; Machado, S.; Budde, H. Systemic Injections of Cannabidiol Enhance Acetylcholine Levels from Basal Forebrain in Rats. *Neurochem. Res.*, **2018**, *43*(8), 1511-1518. [http://dx.doi.org/10.1007/s11064-018-2565-0] [PMID: 29876791]
- [162] Slomski, A. Fewer Seizures With Cannabidiol in Catastrophic Epilepsy. *JAMA*, **2017**, *318*(4), 323. [http://dx.doi.org/10.1001/jama.2017.8846] [PMID: 28742905]
- [163] Devinsky, O.; Patel, A.D.; Cross, J.H.; Villanueva, V.; Wirrell, E.C.; Privitera, M.; Greenwood, S.M.; Roberts, C.; Checketts, D.; VanLandingham, K.E.; Zuberi, S.M. GWPCARE3 Study Group. Effect of Cannabidiol on Drop Seizures in the Lennox-Gastaut Syndrome. *N. Engl. J. Med.*, **2018**, *378*(20), 1888-1897. [http://dx.doi.org/10.1056/NEJMoa1714631] [PMID: 29768152]
- [164] Rubin, R. The Path to the First FDA-Approved Cannabis-Derived Treatment and What Comes Next. *JAMA*, **2018**, *320*(12), 1227-1229. [http://dx.doi.org/10.1001/jama.2018.11914] [PMID: 30193358]
- [165] Ebbert, J.O.; Scharf, E.L.; Hurt, R.T. Medical Cannabis. *Mayo Clin. Proc.*, **2018**, *93*(12), 1842-1847. [http://dx.doi.org/10.1016/j.mayocp.2018.09.005] [PMID: 30522595]
- [166] Freedman, D.A.; Patel, A.D. Inadequate Regulation Contributes to Mislabeled Online Cannabidiol Products. *Pediatr. Neurol. Briefs*, **2018**, *32*, 3. [http://dx.doi.org/10.15844/pedneurbriefs-32-3] [PMID: 30008541]
- [167] Keshavan, M.S.; Giedd, J.; Lau, J.Y.; Lewis, D.A.; Paus, T. Changes in the adolescent brain and the pathophysiology of psychotic disorders. *Lancet Psychiatry*, **2014**, *1*(7), 549-558. [http://dx.doi.org/10.1016/S2215-0366(14)00081-9] [PMID: 26361314]
- [168] Luján, M.A.; Castro-Zavala, A.; Alegre-Zurano, L.; Valverde, O. Repeated Cannabidiol treatment reduces cocaine intake and modulates neural proliferation and CB1R expression in the mouse hippocampus. *Neuropharmacology*, **2018**, *143*, 163-175. [http://dx.doi.org/10.1016/j.neuropharm.2018.09.043] [PMID: 30273593]
- [169] Schiavon, A.P.; Bonato, J.M.; Milani, H.; Guimarães, F.S.; Weffort de Oliveira, R.M. Influence of single and repeated cannabidiol administration on emotional behavior and markers of cell proliferation and neurogenesis in non-stressed mice. *Prog. Neuropsychopharmacol. Biol. Psychiatry*, **2016**, *64*, 27-34. [http://dx.doi.org/10.1016/j.pnpbp.2015.06.017] [PMID: 26187374]
- [170] Carvalhal, R.K.; Souza, M.R.; Santos, M.L.; Guimarães, F.S.; Pobbe, R.L.H.; Andersen, M.L.; Mazaro-Costa, R. Chronic cannabidiol exposure promotes functional impairment in sexual behavior and fertility of male mice. *Reprod. Toxicol.*, **2018**, *81*, 34-40. [http://dx.doi.org/10.1016/j.reprotox.2018.06.013] [PMID: 29936126]
- [171] Iffland, K.; Grotenhermen, F. An Update on Safety and Side Effects of Cannabidiol: A Review of Clinical Data and Relevant Animal Studies. *Cannabis Cannabinoid Res.*, **2017**, *2*(1), 139-154. [http://dx.doi.org/10.1089/can.2016.0034] [PMID: 28861514]
- [172] Sands, T.T.; Rahdari, S.; Oldham, M.S.; Caminha Nunes, E.; Tilton, N.; Cilio, M.R. Long-Term Safety, Tolerability, and Efficacy of Cannabidiol in Children with Refractory Epilepsy: Results from an Expanded Access Program in the US. *CNS Drugs*, **2019**, *33*(1), 47-60. [http://dx.doi.org/10.1007/s40263-018-0589-2] [PMID: 30460546]
- [173] Deutsch, D.G.; Ueda, N.; Yamamoto, S. The fatty acid amide hydrolase (FAAH). *Prostaglandins Leukot. Essent. Fatty Acids*, **2002**, *66*(2-3), 201-210. [http://dx.doi.org/10.1054/plef.2001.0358] [PMID: 12052036]
- [174] Hu, S.S.; Mackie, K. Distribution of the Endocannabinoid System in the Central Nervous System. *Handb. Exp. Pharmacol.*, **2015**, *231*, 59-93. [http://dx.doi.org/10.1007/978-3-319-20825-1\_3] [PMID: 26408158]
- [175] Shin, M.; Ware, T.B.; Lee, H.C.; Hsu, K.L. Lipid-metabolizing serine hydrolases in the mammalian central nervous system: endocannabinoids and beyond. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids*, **2019**, *1864*(6), 907-921. [http://dx.doi.org/10.1016/j.bbalip.2018.08.007] [PMID: 30905349]
- [176] Donvito, G.; Nass, S.R.; Wilkerson, J.L.; Curry, Z.A.; Schurman, L.D.; Kinsey, S.G.; Lichtman, A.H. The Endogenous Cannabinoid System: A Budding Source of Targets for Treating Inflammatory and Neuropathic Pain. *Neuropsychopharmacology*, **2018**, *43*(1), 52-79. [http://dx.doi.org/10.1038/npp.2017.204] [PMID: 28857069]
- [177] Stampanoni Bassi, M.; Gilio, L.; Maffei, P.; Dolcetti, E.; Bruno, A.; Buttari, F.; Centonze, D.; Iezzi, E. Exploiting the Multifaceted Effects of Cannabinoids on Mood to Boost Their Therapeutic Use Against Anxiety and Depression. *Front. Mol. Neurosci.*, **2018**, *11*, 424. [http://dx.doi.org/10.3389/fnmol.2018.00424] [PMID: 30515077]